

BIOAZA 10000 PPM EC

AZADIRACHTIN 10G/L



IRAC (Insecticide Resistance Action Committee) code: 18B.

Description: BIOAZA 10000 PPM EC (Azadirachtin 1 %) is an Insecticide/Nematicide for the control of Aphids, Thrips and Whiteflies on Roses, Aphids, Whiteflies, Leafminers and Spidermites on Tomatoes, Aphids, Cucumber beetles and Thrips on Cucumber, DBM, Aphids, Thrips, Whiteflies and Leafminers on Cabbage and Nematodes on Roses and Cutworms and Nematodes on Tomatoes.

Mixing instructions: Mix BIOAZA 10000 PPM EC by filling the tank halfway with water, adding the required amount while stirring, maintaining agitation, topping up with water, and applying immediately.



ADVANTAGES OF BIOAZA 10000 PPM EC

1. Broad-Spectrum Pest Control

Azadirachtin is effective against a wide range of pests, including insects, mites, and nematodes.

2. Eco-Friendly and Biodegradable

Being a natural compound, Azadirachtin is biodegradable and does not persist in the environment, reducing the risk of pollution.

3. Low Toxicity to Humans and Animals

Azadirachtin has low mammalian toxicity, making it safe for farmers and consumers.

4. Multiple Modes of Action

Azadirachtin acts as an insect growth regulator (IGR), disrupting the molting and reproduction of pests.

5. Reduced Risk of Pest Resistance

Due to its complex mode of action, pests are less likely to develop resistance to Azadirachtin compared to synthetic chemical pesticides.

DISPOSAL

Triple rinse empty container and add washings / rinse off to spray mixture, then perforate or puncture holes on empty containers dispose off in accordance with National Legislation and Regulation for hazard chemical disposal. Do not pour rinsate into any drain or on any site in close proximity to human or animal habitation, water sources, water ways or water bodies. DO NOT reuse empty pesticide containers for any other purpose

AVAILABLE PACKS

50 ml, 100 ml, 250 ml, 500 ml, 1 Litre & 5 Litres

APPLICATION RATE

Crop	Pests	Application Rate, Spray Volume /20L	Timing & Frequency of Application
Roses	False codling moth	2.0L/ ha/ 1000L spray volume of water 40ml/ 20L of water	Apply on the onset of the pests. 3 applications at an interval of 10 days.
Okra	Aphids, Whiteflies, Stink bugs, Leaf miners and Spidermites	1.5L/ ha/ 800L spray volume of water 37.5ml/ 20L of water	Apply on the onset of the pests during vegetative and early flowering stages. 3 applications at an interval of 7 days.
Simsim	Aphids, cutworms, gall midge, spidermites and whiteflies	1.5L/ ha/ 800L spray volume of water 37.5ml/ 20L of water	Apply on the onset of the pests during vegetative and early flowering stages. 3 applications at an interval of 7 days.

NOTICE TO THE USER:

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act (Cap. 346) to use or store a pest control product under unsafe conditions.

SELLERS GUARANTEE:

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.