

FUNGZIL 15 EC

PROCHLORAZ 25 G/L + MYCLOBUTANIL 125 G/L



**FRAC (Fungicide Resistance Action Committee) code: Prochloraz 25 g/L: 3
Myclobutanil 125 g/L: 3**

Description: FUNGZIL 15 EC (Prochloraz 25 g/L + Myclobutanil 125 g/L) is a Fungicide for control of Powdery Mildew on Roses

Mixing instructions: Half fill the spray tank with water. Add the required amount of **FUNGZIL 15 EC** then fill the water to the required level and mix or agitate thoroughly to uniformity. Spray immediately after mixing.



ADVANTAGES OF FUNGZIL 15 EC

1. Provides **strong curative and protective activity** against a wide range of fungal diseases.
2. Combines two DMIs, giving **broader and deeper spectrum control** within the same MOA group.
3. **Enhanced efficacy** on tough pathogens like powdery mildew, scab, and various leaf/fruit diseases.
4. Offers good **systemic movement**, protecting new plant growth.
5. **Improves resistance management** compared to solo use, due to complementary strengths within FRAC 3.

AVAILABLE PACKS

50 g, 100 g, 250 g, 500 g, 1 Kg, 5 Kg & 25Kg

PROCHLORAZ 25 G/L

Category	Information
Site of Action	Inhibition of ergosterol biosynthesis, by inhibition of C14-demethylase (SBI Class I).
Spectrum and Route of Action	Systemic fungicide with protective and eradicant activity.
Uses (Crops & Diseases)	<p>Cereals (wheat, barley, oats, rye): Pseudocercospora spp., Pyrenophora spp., Rhynchosporium spp., Septoria spp., Erysiphe spp.</p> <p>Oilseed Rape (canola): Pyrenopeziza spp., Botrytis spp., Sclerotinia spp.</p> <p>Legumes (peas, beans, lentils): Ascochyta spp., Botrytis spp.</p> <p>Beets: Cercospora spp., Erysiphe spp.</p> <p>Mushrooms: Verticillium fungicola, Mycogone perniciosa</p> <p>Rice: Pyricularia spp. (rice blast)</p> <p>Citrus: Storage diseases (post-harvest fungal infections)</p> <p>Tropical Fruit (mango, papaya, banana, etc.): Storage diseases (post-harvest fungi)</p> <p>Cereals (seed treatment): Cochliobolus spp., Fusarium spp., Pyrenophora spp., Septoria spp.</p> <p>Flax (linseed): Pyrenopeziza spp.</p>

MYCLOBUTANIL 125 G/L

Category	Information
Site of Action	Inhibition of ergosterol biosynthesis, by inhibition of C14-demethylase (SBI Class I).
Spectrum and Route of Action	Systemic fungicide with protective and curative activity, translocated acropetally in the xylem.
Uses (Crops & Diseases)	<p>Vines (grapes): Uncinula necator (powdery mildew)</p> <p>Apples: Leaf scab; powdery mildew</p> <p>Pome Fruit (apple, pear): Powdery mildew; scab</p> <p>Stone Fruit (peach, plum, cherry, apricot): Powdery mildew</p> <p>Cucurbits (cucumber, melon, squash): Powdery mildew</p> <p>Strawberries: Powdery mildew</p> <p>Almonds: Powdery mildew</p> <p>Tomatoes: Powdery mildew</p> <p>Soybeans: Powdery mildew</p> <p>Vegetables (various): Powdery mildew</p> <p>Hops: Powdery mildew</p> <p>Cotton: Powdery mildew</p> <p>Turf & Perennial Grasses: Diseases caused by Ascomycetes, fungi imperfecti, and Basidiomycetes</p> <p>Ornamentals: Powdery mildew and related fungal diseases</p> <p>Barley (seed treatment): Seed- and soil-borne diseases</p> <p>Maize (seed treatment): Seed- and soil-borne diseases</p> <p>Cotton (seed treatment): Seed- and soil-borne diseases</p> <p>Rice (seed treatment): Seed- and soil-borne diseases</p> <p>Wheat (seed treatment): Seed- and soil-borne diseases</p> <p>Bananas (post-harvest): Storage diseases</p>

For complete directions and safety information, please refer to the product label.