

# ZODIAC STAR 30 WDG

AZOXYSTROBIN:200 G/ KG + DIMETHOMORPH:100 G/ KG



FRAC (Fungicide Resistance Action Committee) code: **Azoxystrobin: I I**  
**Dimethomorph: 40**

**Description:** Zodiac Star 30 WDG is used for controlling Powdery Mildew, Downy Mildew and Botrytis on Roses.

**Mixing instructions:** Half-fill the spray tank with water. Add the required amount of **ZODIAC STAR 30 WDG** then fill the water to the required level and mix or agitate thoroughly to uniformity. Spray immediately after mixing.



## ADVANTAGES OF ZODIAC STAR 30 WDG

1. Combines **two complementary modes of action** for strong and reliable disease control.
2. Provides **broad-spectrum activity against oomycetes**, especially downy mildews and late blight.
3. **Preventive, curative, and anti-sporulant effects** reduce disease spread effectively.
4. **Systemic and translaminar movement** ensures protection of new and untreated plant tissues.
5. Enhances **resistance management** by mixing FRAC II and FRAC 40 fungicides.

## AZOXYSTROBIN 200G/ KG

Category	Information
Site of Action	Mitochondrial electron transport inhibition (MET III) by inhibition of ubiquinol oxidase at the Qo (quinone outside) binding site on cytochrome bc1 (complex III).
Spectrum and Route of Action	Systemic fungicide with protective, curative and eradicator activity, absorbed by roots and leaves, translocated acropetally and by translaminar mobility. Inhibits spore germination and mycelial growth and shows antispore activity.
Uses (Crops & Diseases)	<p><b>Cereals:</b> Erysiphe graminis, Puccinia spp., Leptosphaeria nodorum, Septoria tritici, Pyrenophora teres</p> <p><b>Rice:</b> Pyricularia oryzae, Rhizoctonia solani</p> <p><b>Vines (Grapes):</b> Plasmopara viticola, Uncinula necator</p> <p><b>Cucurbits:</b> Sphaerotheca fuliginea, Pseudoperonospora cubensis</p> <p><b>Potatoes &amp; Tomatoes:</b> Phytophthora infestans, Alternaria solani</p> <p><b>Peanuts:</b> Mycosphaerella arachidis, Rhizoctonia solani, Sclerotium rolfsii</p> <p><b>Peaches:</b> Monilinia spp., Cladosporium carpophilum</p> <p><b>Turf:</b> Pythium spp., Rhizoctonia solani</p> <p><b>Bananas:</b> Mycosphaerella spp.</p> <p><b>Pecans:</b> Cladosporium caryigenum</p> <p><b>Citrus:</b> Elsinoë fawcettii, Colletotrichum spp., Guignardia citricarpa</p> <p><b>Coffee:</b> Colletotrichum spp., Hemileia vastatrix</p>

## AVAILABLE PACKS

50 g, 100 g, 250 g, 500 g, 1 Kg, 5 Kg & 25Kg

## DIMETHOMORPH 100G/ KG

Category	Information
Site of Action	
Spectrum and Route of Action	Locally systemic fungicide with protective and antispore activity. Only the (Z)- isomer is intrinsically active, but, because of rapid interconversion of isomers in light, it has no advantage over the (E)- isomer in practice.
Uses (Crops & Diseases)	<p><b>Vines (grapes):</b> Peronosporaceae (downy mildew)</p> <p><b>Potatoes:</b> Phytophthora spp. (late blight)</p> <p><b>Tomatoes:</b> Phytophthora spp. (late blight)</p> <p><b>Other Crops:</b> Oomycete diseases caused by Peronosporaceae and Phytophthora spp.</p>

For complete directions and safety information, please refer to the product label.